

Case Report

Classical imaging in cerebral folate transport deficiency

Sayani Mahal¹, Jaya Pamnani¹ , Manish Parakh² , Sarbesh Tiwari¹ 

¹Department of Diagnostic and Interventional Radiology, All India Institute of Medical Sciences, ²Department of Pediatrics, Dr. S.N. Medical College, Jodhpur, Rajasthan, India.

*Corresponding author:

Sarbesh Tiwari,
Department of Diagnostic
and Interventional Radiology,
All India Institute of Medical
Sciences, Jodhpur, Rajasthan,
India.

sarbesh1984@gmail.com

Received: 15 August 2023
Accepted: 02 September 2023
Epub Ahead of Print: 02 November 2023
Published:

DOI
10.25259/CRCR_138_2023

Quick Response Code:



ABSTRACT

Cerebral folate transport deficiency is an inherited disorder with late infantile-onset caused by nonsense mutations in the folate receptor 1 gene coding for folate receptor alpha. We report a case of a 4-year-old female child, who presented with global neurodevelopmental regression with onset at 15 months of age with an electroencephalogram showing poorly modulated polymorphic delta sleep background. Based on the clinical features, age of onset, and imaging findings, the possibility of cerebral folate transport deficiency (OMIM#613068) was considered. Targeted gene sequencing confirmed the diagnosis.

Keywords: Cerebral folate transport deficiency, Hypomyelination, Neurodevelopmental regression

INTRODUCTION

Cerebral folate transport deficiency is an inherited disorder of brain-specific folate transporter caused by nonsense mutations in the folate receptor 1 gene coding for folate receptor alpha. This leads to the introduction of a premature stop codon in messenger ribonucleic acid causing its rapid degradation.^[1,2] The syndrome is characterized by late infantile onset, severe developmental regression, epilepsy, hypomyelinating leukodystrophy, and low 5-methyltetrahydrofolate (5-MTHF) concentration in the cerebrospinal fluid (CSF) evaluation. Early diagnosis is critical because folinic acid therapy can potentially reverse the clinical symptoms.

CASE REPORT

A 4-year-old female child with non-consanguineous parents presented with global neurodevelopmental regression starting at 15 months of age. She had associated axial myoclonic seizures 6–7 episodes every day, frequent falls, ataxic gait, and language regression. A detailed neurological examination confirmed the global developmental delay, poor eye contact, incomprehensible sounds, unsteady gait, and hypotonia. The fundus evaluation was normal. The thyroid function test, serum folic acid, and Vitamin B12 levels were within normal limits. An electroencephalogram was done, which showed a poorly modulated polymorphic delta sleep background and the absence of any age-appropriate sleep features. There were pseudo-periodic generalized epileptiform discharges. Magnetic resonance imaging (MRI) brain showed diffuse cerebellar atrophy [Figure 1a]. An abnormal T2 hyperintense signal was noted involving the cerebral subcortical white matter on the T2-weighted image [Figure 1b]. The corresponding areas

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, transform, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2023 Published by Scientific Scholar on behalf of Case Reports in Clinical Radiology




Figure 1: Magnetic resonance imaging brain: Axial T2 weighted image (a) shows diffuse cerebellar atrophy. (b) Axial T2-weighted image at the level of the basal ganglia shows abnormal T2-hyperintense signal involving the cerebral subcortical white matter (yellow arrow), which is inappropriate for age. These areas are hypointense on T1 images (c). These features were suggestive of hypomyelination. (d) Axial susceptibility-weighted imaging shows blooming in the subcortical white matter of the bilateral frontal region (yellow arrows), suggesting calcification.

were hypointense on T1 images [Figure 1c]. These features were suggestive of hypomyelination. Axial susceptibility-weighted imaging showed blooming in the subcortical white matter of the bilateral frontal region and basal ganglia, suggesting calcification [Figure 1d]. Computed tomography of the head [Figure 2] confirmed the presence of symmetric calcification in frontal subcortical white matter and basal ganglia.

Based on the clinical features, age of onset, and imaging findings, the possibility of cerebral folate transport deficiency was considered. Targeted gene sequencing was done, which showed a rare homozygous deletion of single base pair (c. 187 del) in exon 3 of the FOLR1 gene (chr11:g.72195289del; Depth: 139x) that results in frame-shift and truncation of the protein 68 amino acids downstream to the codon 63 (p.Asn63MetfsTer68; ENST00000393679.5), confirming the diagnosis.

DISCUSSION

Mutations in the FOLR1 gene result in a brain-specific folate transport defect characterized by a late infantile-onset of severe developmental regression, sometimes commencing in the 2nd or 3rd year of life. Folate transport into cells is regulated by FOLR1, which is a glycoposphatidylinositol-anchored cell membrane protein. Genetic variants have been found to be responsible for different pathologic disorders.^[2-4] Homozygous mutations and compound heterozygous mutations lead to autosomal recessive disorders, while neurodegeneration is caused by homozygous 18-bp in-frame duplication. In this case, we have added a novel homozygous deletion of a single base pair (c. 187 del) variant in the FOLR1 genotype spectrum.

Almost all cases of cerebral folate transporter deficiency demonstrate significantly low levels of 5-MTHF level in CSF.




Figure 2: Computed tomography of the head (blue arrows in a and b) confirmed the presence of symmetric calcification in basal ganglia and frontal subcortical white matter.

These children present with refractory myoclonic epilepsy and ataxia, as seen in our patient.^[1]

Hypomyelination is a well-established association with this condition. Other common central nervous system abnormalities reported in the literature include cerebral atrophy, focal T2 hyperintense white matter lesions, and diffuse hyperintensities in periventricular, and subcortical white matter on T2-weighted images. The focal T2-hyperintense lesion in periventricular and subcortical white matter is more commonly found in older patients and most probably represents areas of gliosis.^[2] Cerebellar atrophy with enlargement of cisterna magna, mild hypoplasia of the vermis, and brainstem atrophy are seen in some cases. Calcification at basal ganglia and the subcortical white matter has been reported in the literature and was also seen in our case.^[1] Imaging differentials of subcortical white matter signal changes with calcification include Kearns–Sayre syndrome, Cockayne syndrome, and Aicardi–Goutieres syndrome. The absence of progressive external ophthalmoplegia and

cardiac conduction defects were against the diagnosis of Kearns–Sayre syndrome. In Aicardi–Goutières syndrome, calcifications are more frequently punctuate and distributed in the basal ganglia, the deep and periventricular white matter^[5] Cockayne syndrome is characterized by cutaneous photosensitivity, significant white matter loss, and ventricular dilatation, which were not present in our case.^[6,7]

Cerebral folate transport deficiency is not exclusively caused by FOLR1 genetic variants. Hereditary folate malabsorption, MTHF, and dihydrofolate reductase deficiency are other significant causes. Normal red blood cell count and plasma concentration of homocysteine and folate within normal limits exclude all these conditions.^[2]

It has been reported in several studies that combined oral and intravenous oral folinic acid can improve both clinical symptoms and white matter abnormalities.

CONCLUSION

Cerebral folate deficiency must be kept as differential in central hypomyelination and calcification. This being a potentially treatable inborn error of metabolism, a prompt diagnosis and early initiation of treatment is important to ensure a better outcome.

TEACHING POINTS

1. Cerebral folate transport deficiency is an inherited neurodegenerative autosomal recessive disorder resulting from brain-specific folate transporter deficiency early in life.
2. Diagnosis is based on the clinical phenotype of severe developmental regression, epilepsy, progressive ataxia, and reduced 5-methyltetrahydrofolate in CSF analysis.
3. Magnetic Resonance Imaging (MRI) findings of central hypomyelination with basal ganglia and subcortical calcification in appropriate clinical scenarios should raise a suspicion of cerebral folate transport deficiency.

MCQs

1. Most common age of onset of cerebral folate transporter deficiency is-
 - a. Neonatal
 - b. Adult onset
 - c. Late infantile
 - d. Adolescence

Answer Key: c

2. A substrate that is affected in CSF in cases of cerebral folate transporter deficiency is-
 - a. Neuregulin 1
 - b. P- glycoprotein

- c. Contactin 2
- d. 5-methyltetrahydrofolate (5-MTHF)

Answer Key: d

3. The most common clinical feature of cerebral folate transporter deficiency is -
 - a. Neuroregression
 - b. Refractory epilepsy
 - c. Headache
 - d. Reduced visual acuity

Answer Key: b

4. Which of the following is not a differential diagnosis of cerebral folate transporter deficiency?
 - a. Kearns–Sayre syndrome
 - b. Cockayne syndrome
 - c. Sturge–Weber syndrome
 - d. Aicardi–Goutieres syndrome

Answer Key: c

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Use of artificial intelligence (AI)-assisted technology for manuscript preparation

The author(s) confirms that there was no use of artificial intelligence (AI)-assisted technology for assisting in the writing or editing of the manuscript and no images were manipulated using AI.

REFERENCES

1. Brunetti S, Malerba L, Giordano L, Parrini E, Guerrini R, Palumbo G, *et al.* Cerebral folate transporter deficiency syndrome in three siblings: Why genetic testing for developmental and epileptic encephalopathies should be performed early and include the FOLR1 gene. *Am J Med Genet A* 2021;185:2526-31.
2. Grapp M, Just IA, Linnankivi T, Wolf P, Lücke T, Häusler M, *et al.* Molecular characterization of folate receptor 1 mutations delineates cerebral folate transport deficiency. *Brain* 2012;135:2022-31.
3. Pérez-Dueñas B, Toma C, Ormazábal A, Muchart J, Sanmartí F, Bombau G, *et al.* Progressive ataxia and myoclonic epilepsy in a patient with a homozygous mutation in the FOLR1 gene. *J Inher Metab Dis* 2010;33:795-802.

4. Pope S, Artuch R, Heales S, Rahman S. Cerebral folate deficiency: Analytical tests and differential diagnosis. *J Inherit Metab Dis* 2019;42:655-72.
5. Uggetti C, La Piana R, Orcesi S, Egitto MG, Crow YJ, Fazzi E. Aicardi-Goutieres syndrome: Neuroradiologic findings and follow-up. *Am J Neuroradiol* 2009;30:1971-6.
6. Koob M, Laugel V, Durand M, Fothergill H, Dalloz C, Sauvanaud F, *et al.* Neuroimaging in cockayne syndrome. *Am J Neuroradiol* 2010;31:1623-30.
7. Gonçalves FG, Caschera L, Teixeira SR, Viaene AN, Pinelli L, Manka K, *et al.* Intracranial calcifications in childhood: Part 2. *Pediatr Radiol* 2020;50:1448-75.

How to cite this article: Mahal S, Pamnani J, Parakh M, Tiwari S. Classical imaging in cerebral folate transport deficiency. *Case Rep Clin Radiol*, doi: 10.25259/CRCR_138_2023